

#### 4. Publications

- Awaya et al. (2017) Evaluating the differences in modeling biophysical attributes between deciduous broadleaved and evergreen conifer forests using low-density small-footprint LiDAR data. *Remote Sensing* **9**: 572, doi: :10.3390/rs9060572.
- Ballav et al. (2012) Simulation of CO<sub>2</sub> concentration over East Asia using the regional transport model WRF-CO<sub>2</sub>. *Journal of the Meteorological Society of Japan* **90**: 959-976.
- Cao et al. (2019) Nitrogen deposition and responses of forest structure to nitrogen deposition in a cool-temperate deciduous forest. *Forests* **10**: 631, doi:10.3390/f10080631.
- Chen et al. (2017) Dissolved organic carbon (DOC) input to the soil: DOC fluxes and their partitions during the growing season in a cool-temperate broad-leaved deciduous forest, central Japan. *Ecological Research* **32**: 713-724.
- He et al. (2013) Evaluation and improvement of MODIS gross primary productivity in typical forest ecosystems of East Asia based on eddy covariance measurements. *Journal of Forest Research* **18**: 31-40.
- He et al. (2021) Microbial seasonality promotes soil respiratory carbon emission in natural ecosystems: A modeling study. *Global Change Biology* **27**: 3035-3051.
- Hirata et al. (2008) Spatial distribution of carbon balance in forest ecosystems across East Asia. *Agricultural and Forest Meteorology* **148**: 761-775.
- Ichii et al (2013) Site-level model-data synthesis of terrestrial carbon fluxes in the CarboEastAsia eddy-covariance observation network: Toward future modeling efforts. *Journal of Forest Research* **18**: 13-20.
- Ichii et al. (2010) Multi-model analysis of terrestrial carbon cycles in Japan: Limitations and implications of model calibration using eddy flux observations. *Biogeosciences* **7**: 2061-2081.
- Inatomi et al. (2010) Greenhouse gas budget of a cool-temperate deciduous broad-leaved forest in Japan estimated using a process-based model. *Ecosystems* **13**: 472-483.
- Inoue et al. (2014) Detection of the different characteristics of year-to-year variation in foliage phenology among deciduous broad-leaved tree species by using daily continuous canopy surface images. *Ecological Informatics* **22**: 58-68.
- Ishidoya et al. (2013) O<sub>2</sub>:CO<sub>2</sub> exchange ratios observed in a cool temperate deciduous forest ecosystem of central Japan, *Tellus* **65B**: 21120, doi.org/10.3402/tellusb.v65i0.21120.
- Ishidoya et al. (2015) Observation of O<sub>2</sub>:CO<sub>2</sub> exchange ratio for net turbulent fluxes and its application to forest carbon cycles. *Ecological Research* **30**: 225-234.
- Ito (2010) Changing ecophysiological processes and carbon budget in East Asian ecosystems under near-future changes in climate: implications for long-term monitoring from a process-based model. *Journal of Plant Research* **123**: 577-588.
- Ito (2010) Evaluation of defoliation impacts of tropical cyclones on the forest carbon budget using flux data and a process-based model. *Journal of Geophysical Research* **115**: G04013 doi:10.1029/2010JG001314.
- Ito et al. (2006) Seasonal variation in leaf properties and ecosystem carbon budget in a cool-temperate deciduous broad-leaved forest: simulation analysis at Takayama site, Japan. *Ecological Research* **21**:137-149.
- Ito et al. (2007) Examination of model-estimated ecosystem respiration by use of flux measurement data from a cool-temperate deciduous broad-leaved forest in central Japan. *Tellus* **59B**, 616-624.
- Ito et al. (2015) Synergies between observational and modeling studies at the Takayama site: toward a better understanding of terrestrial ecosystem processes. *Ecological Research* **30**: 201-210.
- Jang et al. (2010) Mapping evapotranspiration using MODIS and MM5 four-dimensional data assimilation. *Remote Sensing of Environment* **114**: 657-673.

- Jia et al. (2003) Temporal and spatial variability of soil respiration in a cool temperate broad-leaved forest. 1. Measurement of spatial variance and factor analysis. *Japanese Journal of Ecology* **53**: 13-22. (in Japanese with English abstract)
- Kamakura et al. (2011) Simulations and observations of patchy stomatal behavior in leaves of *Quercus crispula*, a cool-temperate deciduous broad-leaved tree species. *Journal of Plant Research* **125**: 339-349.
- Kishimoto-Mo et al. (2015) Contribution of soil moisture to seasonal and annual variations of soil CO<sub>2</sub> efflux in a humid cool-temperate oak-birch forest in central Japan. *Ecological Research* **30**: 311-325.
- Kondo et al. (2001) A numerical simulation of the daily variation of CO<sub>2</sub> in the central part of Japan -summer case-. *Journal of Meteorological Society of Japan* **79**: 11-21.
- Kondo et al. (2005) Refixation of respired CO<sub>2</sub> by understory vegetation in a cool-temperate deciduous forest in Japan. *Agricultural and Forest Meteorology* **134**: 110-121.
- Kondo et al. (2010) Radiocarbon-based residence time estimates of soil organic carbon in a temperate forest: Case study for the density fractionation for Japanese volcanic ash soil. *Nuclear Instruments and Methods in Physics Research B* **268**: 1073-1076.
- Kondo et al. (2015) Impact of anomalous climates on carbon allocation to biomass production of leaves, woody components, and fine roots in a cool-temperate deciduous forest. *Agricultural and Forest Meteorology* **201**: 38-50.
- Kume et al. (2011) The ratio of transmitted near-infrared radiation to photosynthetically active radiation (PAR) increases in proportion to the adsorbed PAR in the canopy. *Journal of Plant Research* **124**: 99-106.
- Lee et al. (2002) Effects of rainfall events on soil CO<sub>2</sub> flux in a cool temperate deciduous broad-leaved forest. *Ecological Research* **17**: 401-409.
- Lee et al. (2003) Seasonal changes in the contribution of root respiration to total soil respiration in a cool-temperate deciduous forest. *Plant and Soil* **255**: 311-318.
- Lee et al. (2005) The importance of root respiration in annual soil carbon fluxes in a cool-temperate deciduous forest. *Agricultural and Forest Meteorology* **134**: 95-101.
- Lee et al. (2006) Soil respiration of forest ecosystems in Japan and global implications. *Ecological Research* **21**: 828-839.
- Mariko et al. (2000) Measurement of CO<sub>2</sub> fluxes from soil and snow surfaces with open dynamic chamber technique. *Environmental Sciences* **13**: 69-74.
- Mariko et al. (2000) Winter CO<sub>2</sub> flux from soil and snow surfaces in a cool-temperate deciduous forest, Japan. *Ecological Research* **15**: 363-372.
- Mikami et al. (2006) Automatic detection of forest canopy gaps and estimation of leaf area index (LAI) using the digital fish-eye camera's images. *Journal of the Japan Society of Photogrammetry and Remote Sensing* **45**: 13-22. (in Japanese with English abstract)
- Mo et al. (2005) Interannual variation in CO<sub>2</sub> effluxes from soil and snow surfaces in a cool-Temperate deciduous broad-leaved forest. *Phyton* **45**: 99-107.
- Mo et al. (2005) Seasonal and annual variations in soil respiration in a cool-temperate deciduous broad-leaved forest in Japan. *Agricultural and Forest Meteorology* **134**: 81-94.
- Morozumi et al. (2023) Contributions of the understory and midstory to total canopy solar-induced chlorophyll fluorescence in a ground-based study in conjunction with seasonal gross primary productivity in a cool-temperate deciduous broadleaf forest. *Remote Sensing of Environment* **284**: 113340, doi: 10.1016/j.rse.2022.113340.
- Motohka et al. (2010) Applicability of Green-Red Vegetation Index for remote sensing of vegetation phenology. *Remote Sensing* **2**: 2369-2387.
- Muraoka & Koizumi (2005) Photosynthetic and structural characteristics of canopy and shrub trees in a cool-temperate deciduous forest: implication to the ecosystem carbon gain. *Agricultural and Forest Meteorology* **134**: 39-59.
- Muraoka & Koizumi (2006) Leaf and shoot ecophysiological properties and their role in photosynthetic carbon gain of cool-temperate deciduous forest trees. In: Kawatata H and Awaya Y (eds.) *Global climate*

*change and response of carbon cycle in the Equatorial Pacific and Indian Oceans and adjacent landmasses*. Elsevier Oceanography series vol. 73, Elsevier, pp.417-443.

- Muraoka & Koizumi (2009) "Satellite Ecology" for linking ecology, remote sensing and micrometeorology from plot to regional scales for ecosystem structure and function study. *Journal of Plant Research* **122**: 3-20.
- Muraoka (2022) Chapter 1: Phenology of photosynthesis in a deciduous broadleaf forest: implications for the carbon cycle in a changing environment. In: Li F. Awaya Y., Kageyama K., Wei Y. (eds.) *River basin environment: evaluation, management and conservation*, Springer Nature Singapore, doi:10.1007/978-981-19-4070-5\_1.
- Muraoka et al. (2010) Effects of seasonal and interannual variation in leaf photosynthesis and canopy leaf area index on gross primary production in a cool-temperate deciduous broadleaf forest in Takayama, Japan. *Journal of Plant Research* **123**: 563-576.
- Muraoka et al. (2013) Spectral vegetation indices as the indicator of canopy photosynthetic productivity in a deciduous broadleaf forest. *Journal of Plant Ecology* **6**: 393-407.
- Muraoka et al. (2015) [Editorial] Long-term and interdisciplinary research on forest ecosystem functions: challenges at Takayama site since 1993. *Ecological Research* **30**: 197-200.
- Murayama et al. (2003) Temporal variations of atmospheric CO<sub>2</sub> concentration in a temperate deciduous forest in central Japan. *Tellus* **55B**: 232-243.
- Murayama et al. (2005) Statistical analyses of inter-annual variations in the vertical profile of atmospheric CO<sub>2</sub> concentration and carbon budget in a cool-temperate deciduous forest in Japan. *Agricultural and Forest Meteorology* **134**: 17-26.
- Murayama et al. (2010) Seasonal variations of atmospheric CO<sub>2</sub>, δ<sup>13</sup>C, and δ<sup>18</sup>O at a cool temperate deciduous forest in Japan: Influence of Asian monsoon. *Journal of Geophysical Research* **115**: D17304, doi:10.1029/2009JD013626.
- Musavi et al. (2017) Stand age and species richness dampen interannual variation of ecosystem-level photosynthetic capacity. *Nature ecology & evolution* **1**: e0048, doi:10.1038/s41559-016-0048.
- Nagai & Nasahara (2017) Seasonal leaf phenology data for 12 tree species in a cool-temperate deciduous broadleaved forest in Japan from 2005 to 2014. *Ecological Research* **32**:107-107.
- Nagai et al. (2010) Field experiments to test the use of the normalized difference vegetation index for phenology detection. *Agricultural and Forest Meteorology* **150**: 152-160.
- Nagai et al. (2010) What makes the satellite-based EVI-GPP relationship unclear in a deciduous broad-leaved forest? *Ecological Research* **25**: 359-365.
- Nagai et al. (2011) The necessity and availability of noise-free daily satellite-observed NDVI during rapid phenological changes in terrestrial ecosystems in East Asia. *Forest Science and Technology* **7**: 174-183.
- Nagai et al. (2011) Using digital camera images to detect canopy condition of deciduous broad-leaved trees. *Plant Ecology and Diversity* **4**: 78-88.
- Nagai et al. (2013) Detection of bio-meteorological year-to-year variation by using digital canopy surface images of a deciduous broad-leaved forest. *SOLA* **9**: 106-110.
- Nagai et al. (2013) Utility of information in photographs taken upwards from the floor of closed-canopy deciduous broadleaved and closed-canopy evergreen coniferous forests for continuous observation of canopy phenology. *Ecological Informatics* **18**: 10-19.
- Nagai et al. (2014) Relationship between spatio-temporal characteristics of leaf-fall phenology and seasonal variations in near surface- and satellite-observed vegetation indices in a cool-temperate deciduous broad-leaved forest in Japan. *International Journal of Remote Sensing* **35**: 3520-3536.
- Nagai et al. (2014) Year-to-year blooming phenology observation by using time-lapse digital camera images. *Journal of Agricultural Meteorology* **70**: 163-170.
- Nagai et al. (2015) Uncertainties involved in leaf fall phenology detected by digital camera. *Ecological Informatics* **30**: 124-132.
- Nagai et al. (2016) Review: advances in in situ and satellite phenological observations in Japan. *International Journal of Biometeorology* **60**: 615-627.
- Nagai et al. (2017) Seasonality of leaf litter and leaf area index data for various tree species in a cool-temperate deciduous broad-leaved forest, Japan, 2005-2014. *Ecological Research* **32**: 297.
- Nagai et al. (2018) 8 million phenological and sky images from 29 ecosystems from the Arctic to the tropics: the Phenological Eyes Network. *Ecological Research* **33**: 1091-1092.
- Nakashima et al. (2021) Area-ratio Fraunhofer line depth (aFLD) method approach to estimate solar-induced chlorophyll fluorescence in low spectral resolution spectra in a cool-temperate deciduous broadleaf forest. *Journal of Plant Research* **134**:713-728.
- Nasahara & Nagai (2015) Development of an in-situ observation network for terrestrial ecological remote

- sensing — the Phenological Eyes Network (PEN). *Ecological Research* **30**: 211-223.
- Nasahara et al. (2008) Vertical integration of leaf area index in a Japanese deciduous broad-leaved forest. *Agricultural and Forest Meteorology* **148**: 1136-1146.
- Nishimura et al. (2004) Evaluation of carbon budgets of a forest floor *Sasa senanensis* community in a cool-temperate forest ecosystem, central Japan. *Japanese Journal of Ecology* **54**: 143-158. (in Japanese with English abstract)
- Noda et al. (2015) Phenology of leaf morphological, photosynthetic, and nitrogen use characteristics of canopy trees in a cool-temperate deciduous broadleaf forest at Takayama, central Japan. *Ecological Research* **30**: 247-266.
- Noda et al. (2021) Phenology of leaf optical properties and their relationship to mesophyll development in cool-temperate deciduous broad-leaf trees. *Agricultural and Forest Meteorology* **297**: 108236, doi: 10.1016/j.agrformet.2020.108236.
- Noguchi et al. (2007) Biomass and production of fine roots in Japanese forests. *Journal of Forest Research* **12**: 83-95.
- Noh et al. (2016) Responses of soil, heterotrophic, and autotrophic respiration to experimental open-field soil warming in a cool-temperate deciduous forest. *Ecosystems* **19**: 504-520.
- Noh et al. (2017) Different responses of soil, heterotrophic and autotrophic respirations to a 4-year soil warming experiment in a cool-temperate deciduous broadleaved forest in central Japan. *Agricultural and Forest Meteorology* **247**: 560-570.
- Ohtsuka et al. (2005) Biometric based estimates of net primary production (NPP) in a cool-temperate deciduous forest stand beneath a flux tower. *Agricultural and Forest Meteorology* **134**: 27-38.
- Ohtsuka et al. (2009) On linking multiyear biometric measurements of tree growth with eddy covariance-based net ecosystem production. *Global Change Biology* **15**: 1015-1024.
- Ohtsuka et al. (2014) Role of coarse woody debris in the carbon cycle of Takayama forest, central Japan. *Ecological Research* **29**: 91-101.
- Piao et al. (2011) Contribution of climate change and rising CO<sub>2</sub> to terrestrial carbon balance in East Asia: A multi-model analysis. *Global and Planetary Change* **75**: 133-142.
- Saigusa et al. (2002) Gross primary production and net ecosystem production of a cool-temperate deciduous forest estimated by the eddy covariance method. *Agricultural and Forest Meteorology* **112**: 203-215.
- Saigusa et al. (2005) Inter-annual variability of carbon budget components in an AsiaFlux forest site estimated by long-term flux measurements. *Agricultural and Forest Meteorology* **134**: 4-16.
- Saigusa et al. (2005) Inter-annual variability of carbon budget components in a cool-temperate deciduous forest in Japan (Takayama, AsiaFlux). *Phyton* **45**: 81-88.
- Saigusa et al. (2008) Temporal and spatial variations in the seasonal patterns of CO<sub>2</sub> flux in boreal, temperate, and tropical forests in East Asia. *Agricultural and Forest Meteorology* **148**: 700-713
- Saigusa et al. (2010) Impact of meteorological anomalies in the 2003 summer on gross primary productivity in East Asia, *Biogeosciences* **7**: 641-655.
- Saigusa et al. (2013) Dataset of CarboEastAsia and the uncertainties in the CO<sub>2</sub> budget evaluation caused by different data-processing. *Journal of Forest Research* **18**: 41-48.
- Saitoh et al. (2012) Assessing the use of camera-based indices for characterizing canopy phenology in relation to gross primary production in a deciduous broad-leaved and an evergreen coniferous forest in Japan. *Ecological Informatics* **11**: 45-54.
- Saitoh et al. (2012) Examination of the extinction coefficient in the Beer-Lambert law for an accurate estimation of the forest canopy leaf area index. *Forest Science and Technology* **8**: 67-76.
- Saitoh et al. (2012) Functional consequences of differences in canopy phenology for the carbon budgets of two cool-temperate forest types: simulations using the NCAR/LSM model and validation using tower flux and biometric data. *Eurasian Journal of Forest Research* **15**: 19-30.
- Saitoh et al. (2015) Effects of canopy phenology on deciduous overstory and evergreen understory carbon budgets in a cool-temperate forest ecosystem under ongoing climate change. *Ecological Research* **30**: 267-277.
- Saitoh et al. (2022) Chapter 3: Forest carbon sequestration in mountainous region in Japan under ongoing climate change: implication for future research. In: Li F. Awaya Y., Kageyama K., Wei Y. (eds.) *River basin environment: evaluation, management and conservation*, Springer Nature Singapore, doi: 10.1007/978-981-19-4070-5\_3.
- Sakai & Akiyama (2005) Quantifying the spatio-temporal variability of net primary production of the understory species, *Sasa senanensis*, using multipoint measuring techniques. *Agricultural and Forest Meteorology* **134**: 60-69.
- Sakai et al. (2005) Microsite variation in light availability and photosynthesis in a cool-temperate deciduous broadleaf forest in central Japan. *Ecological Research* **20**: 537-545.

- Sakai et al. (2006) The contribution of gross primary production of understory dwarf bamboo, *Sasa senanensis*, in a cool-temperate deciduous broadleaf forest in central Japan. *Forest Ecology and Management* **236**: 259-267.
- Sakai et al. (2009) Quantitative analysis of canopy photosynthesis influenced by light simulation models. In: Jason D. Creighton and Paul J. Roney (eds.) *Forest Canopies: Forest Production, Ecosystem Health, and Climate Conditions*, Nova Science Publishers, NY, USA, pp.51-70.
- Sakurai et al. (2015) Inversely estimating the vertical profile of the soil CO<sub>2</sub> production rate in a deciduous broadleaf forest using a particle filtering method. *PLoS ONE* **10**: e0119001.
- Satomura et al. (2006) Seasonal patterns of fine root demography in a cool-temperate deciduous forest in central Japan. *Ecological Research* **21**: 741-753.
- Shen et al. (2020) Climate responses of ring widths and radial growth phenology of *Betula ermanii*, *Fagus crenata* and *Quercus crispula* in a cool temperate forest in central Japan. *Trees* **34**: 679-692.
- Shen et al. (2022) Effects of climate on the tree ring density and weight of *Betula ermanii* in a cool temperate forest in central Japan. *Trees* **36**: 1597-1605.
- Suminokura et al. (2018) Non-destructive measurement of soil respiration in a grassland ecosystem using the multiple-microchambers method. *Ecological Research* **33**: 471-477.
- Uchida et al. (2005) Microbial activity and litter decomposition under snow cover in a cool-temperate broad-leaved deciduous forest. *Agricultural and Forest Meteorology* **134**: 102-109.
- Uchida et al. (2008) Ecosystem-scale carbon isotope ratio of respired CO<sub>2</sub> in cool-temperate deciduous forests under Asian Monsoon climate. *Journal of Geophysical Research* **113**, G02015, doi:10.1029/2007JG000574.
- Yamaji et al. (2008) Scaling-up technique for net ecosystem productivity of deciduous broadleaved forests in Japan using MODIS data. *Ecological Research* **23**, 765-775.
- Yamamoto et al. (1999) Seasonal and inter-annual variation of CO<sub>2</sub> flux between a temperate forest and the atmosphere in Japan. *Tellus* **51B**: 402-413.
- Yamamoto et al. (2022) Uncertainty quantification in land surface temperature retrieved from Himawari-8/AHI data by operational algorithms. *ISPRS Journal of Photogrammetry and Remote Sensing* **191**: 171-187.
- Yamamoto et al. (2023) Detection of vegetation drying signals using diurnal variation of land surface temperature: Application to the 2018 East Asia heatwave. *Remote Sensing of Environment* **291**: 113572, doi: 10.1016/j.rse.2023.113572.
- Yamamoto et al. (2013) Research on the carbon cycle in the forest ecosystems -Twenty years observation of the carbon exchange at Takayama site and Its Extension to the AsiaFlux Network-. *Tenki*, **60**: 359-370 (in Japanese).
- Yashiro et al. (2012) The effect of dense dwarf bamboo (*Sasa senanensis*) understory on soil respiration before and after clearcutting of cool temperate deciduous broad-leaved forest. *Ecological Research* **27**: 577-586.
- Yonemura et al. (2013) Vertical soil-air CO<sub>2</sub> dynamics at the Takayama deciduous broadleaved forest AsiaFlux site. *Journal of Forest Research* **18**: 49-59.

#### AsiaFlux Newsletter Articles

[No.8] Synthesis Workshop on the Carbon Budget in Asian Monitoring Network -The decennial anniversary of the observation at Takayama site-

[No.17] Special Issue on the Long-term Carbon Exchange at Takayama Forest Site, Japan by N. SAIGUSA et al.

[No.20] Validation of Terrestrial Carbon Cycle Model with AsiaFlux Data by A. Ito